

Table 1. **REQUIREMENTS FOR ROUTINE VACCINATION OF IMMIGRANTS EXAMINED OVERSEAS WHO ARE NOT FULLY VACCINATED OR HAVE NO DOCUMENTATION.** (All vaccines may be given at the same time, but at different sites on the body.)

| Vaccine | Age | | | | | | |
|--------------------------|------------------------------|---|-------------------------|-----------|------------|--|-------------------|
| | Birth-1 month | 2-11 months | 12 months-4 years | 5-6 years | 7-17 years | 18-64 years | 65 years or older |
| DTP/DTaP; may include DT | NO | YES | | | NO | | |
| Td | NO | | | | YES | | |
| Polio; IPV or OPV | NO | YES | | | | NO | |
| MMR | NO | | YES, if born after 1956 | | | | NO |
| Hib | NO | YES | | NO | | | |
| Hepatitis B | Yes, through 19 years of age | | | | | NO | |
| Varicella | NO | | YES | | | | |
| Pneumococcal | NO | YES, through 23 months of age (for PCV) | | NO | | | YES (for PPV) |
| Influenza | NO | | | | | YES, 50 years or older (annually, each flu season) | |

DTP=diphtheria and tetanus toxoids and pertussis vaccine; DTaP=diphtheria and tetanus toxoids and acellular pertussis vaccine; DT=pediatric formulation diphtheria and tetanus toxoids; Td=adult formulation tetanus and diphtheria toxoids; IPV=inactivated polio vaccine (killed); OPV=oral polio vaccine (live); MMR=combined measles, mumps, rubella vaccine; Hib=*Haemophilus influenzae* type b conjugate vaccine; PCV=pneumococcal conjugate vaccine; PPV= pneumococcal polysaccharide vaccine.